

令和6年6月 第93巻 第6号 (第1077号)

「巻頭	言」上下水道一体での事業推進に向けて		··松	原		誠•••	(1)
			仲	田	義孝	信	
「報文」	におい嗅ぎ GC を用いた水道水における珪液 樹脂臭に関する調査と浄水処理の改善	上藻由来の	·阿	部	季晃和	欣 文···	(2)
	関加矢に関する調査と伊小処理の以音		新二本	井	和秀	也治	
			渡	辺	直	人	
「報文」	有機物の水質検査に用いる採水容器として 適用可能性の検討		川	本	佳	澄	
			三関	浦川	佳晃慎	也	
			荒中	井	活	人	
		ての樹脂容器の	屮 :: 金	井敷	喬	彦 毅···	(13)
	週用り配性の快割		長名	-	哲	批	(=0)
			立木	石村	哲浩美	之紀美	
			藤小	井坂	裕浩	美司	
			小小	林	憲	弘	
「資料」	令和6年度水道施設関係予算の概要	·国土交通省水管理·国土的	呆全局	引水道	事業	[課…	(22)
「資料」	令和6年度地方財政の見通しと水道事業・・	総務省自治財政局	哥公営	企業	経営	室…	(25)
「資料」	令和6年度独立行政法人水資源機構予算の)概要					
							(54)
文 献	抄 録			• • • • •	• • • • •	•••••	(60)
	tの観点から見た水道料金滞納手数料と財政支持					睦…	(60)
持続可 Filton	「能な水管理のための都市水資源の回収の評価 飛行場のケーススタディ (イギリス)	:	····/]\	林	尚	貴…	(62)
ペルフ 粒状活	フルオロアルキル化合物およびポリフルオロア) 5性炭(GACs)の再生および賦活化に関する研	レキル化合物(PFAS)を含 f究の現状	^Ե <u>.</u> Է	原	佳	奈…	(64)
浄水処	L理におけるジェオスミン及び2- メチルイソボ	ルネオールの除去・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・	····Ξ	好	太	郎…	(66)
文 献	目 録			• • • • •			(69)
新聞情	報目録						(71)
ニューフ	、(巻頭)	「お知らせ」本協会共催・	協賛	・後抱	その行	· 事····	(80)
	、り・・・・・・・・・・・・・・・・(巻頭)	「お知らせ」今後開催予算					
	日本水道協会第104回総会の (巻頭) 開催について	「会告」 日本水道協会「オ 協力専門家登録制		ニア	国際. 案内		(87)
「会告」	令和6年度 日本水道協会主要行事予定表(巻頭)	「会告」日本水道協会 「研修講師登録制	度」の	りご案			(88)
「会告」	令和6年度 日本水道協会研修会開催日程案 ······(巻頭)	日本水坦肠会発行凶青日新	永				(90)
「会告」	公益社団法人日本水道協会 受信力・情報発信力の強化に向けて・・・(巻頭)	JWWA(日本水道協会) 「会告」法律・経営無料木	現格 目	録・・・	 л .		(92) (04)
		「お知らせ」今月の新蔵書	ロロバマノ 書•・・・・	······			(95)
お知り	らせ」 水道協会雑誌・水道研究発表会 講演集掲載論文等の J-ST AGE ・・・ (巻頭)	水道協会雑誌投稿規程					
□ Merulat 1	への公開について	会誌編集委員会及び抄録者					
	水道用品検査実績(令和6年3月分)····(74) JIS 製品認証事業の認証······(79)	編集後記	• • • • • •	• • • • •	• • • • •	• • • • • •	(100)
141	J10 丞叫心叫于木 / 10 10 10 (19)						

Journal of Japan Water Works Association

Vol. 93 No.6 June 2024

Contents

Implementing the Integrated Approaches to Water and Wastewater Management · · ·
by Makoto MATSUBARA ··· (1)
Investigation of a Resin-like Odorous Compound Found in Drinking Water Derived from Diatoms Using GC-Olfactometry, and Subsequent Improvement of Water Purification Processes · · ·
A resin-like odor observed in tap water from Kawasaki City was investigated using gas chromatography with an odor-sniffing technique. The study estimated that 1-Hexen-3-one is the causative agent, which originates from diatoms such as the genus Aulacoseira found in raw water at the water purification plant. 1-Hexen-3-one was detected in a sample in which Aulacoseira granulata var. angustissima was mono-cultured from raw water and subsequently the cells were disrupted by ultrasonic waves. Furthermore, the concentration of 1-Hexen-3-one increased due to chlorination. Therefore, it was decided to develop a test method to determine the formation potential of 1-Hexen-3-one as an indicator for monitoring raw water. Regarding the control of odor, 1-Hexen-3-one was confirmed to be removed by powdered activated carbon. It should be noted that 1-Hexen-3-one is likely to be removed equivalently or less compared with 2-methyl-isoborneol.
Investigation of Applicability of Plastic Bottle as Water Sampling Container for Organic Substances in Water Quality Test \cdots
by Naoto WATANABE, Kasumi KAWAMOTO, Kouichi MIURA, Shinya SEKIKAWA, Katsuhito ARAI,
Takahiko NAKAI, Takeshi KANASHIKI, Tetsuya HASEGAWA, Hiroyuki TATEISHI, Miki KIMURA,
Hiromi FUJII, Koji KOSAKA and Norihiro KOBAYASHI · · · (13)
We investigated whether plastic bottles could be used as water sampling containers for water quality test items where glass water sampling containers are specified in the testing method of organic substance. In a survey for water utilities, although some water utilities expressed concerns about adsorption and durability, approximately 80% of water utilities wished the use of plastic bottles, indicating that there is a strong need. For the investigation, we used high-density polyethylene (HDPE) bottle, which have high availability and chemical resistance and are used as water sampling containers for water quality testing for inorganic substances. In a comparative test using samples in which the target substance was added to tap water, we confirmed that TOC, formaldehyde, phenols, and haloacetic acids were comparable to glass bottles. However, in a carryover test targeting formaldehyde and phenols, carryover was observed in phenols when HDPE bottles were used. Additionally, during the investigation, we confirmed that the acetone cleaning specified in the testing methods for formaldehyde and phenols is not necessarily necessary when using glass bottles. Based on the above results, we made a proposal to the Ministry of Health, Labor and Welfare's Water Quality Testing Method Review Committee to revise the testing methods.
Abstracts of Foreign References ······(60)